Treating your infection



Your infection	Usually lasts	How to treat yourself better for these infections, now and next time	When should you get help: Contact your GP practice or contact NHS 111 (England), NHS 24 (Scotland of 111), or NHS Direct (Wales dial 0845 4647)
Middle-ear infection	4 days	Have plenty of rest. Drink enough fluids to avoid feeling thirsty. Ask your local pharmacist to recommend medicines to help your symptoms or pain (or both). Fever is a sign the body is fighting the infection and usually gets better by itself in most cases. You can use paracetamol (or ibuprofen) if you or your child are uncomfortable as a result of a fever. Other things you can do suggested by GP or nurse:	 to 8. are possible signs of serious illness and should be assessed urgently. Phone for advice if you are not sure how urgent the symptoms are. If you develop a severe headache and are sick. If your skin is very cold or has a strange colour, or you develop an unusual rash. If you feel confused or have slurred speech or are very drowsy. If you have difficulty breathing. Signs can include: breathing quickly turning blue around the lips and the skin below the mouth skin between or above the ribs getting sucked or pulled in with every breath. If you develop chest pain. If you have difficulty swallowing or are drooling. If you cough up blood. If you are feeling a lot worse. Less serious signs that can usually wait until the next available GP appointment: If you are not improving by the time given in the 'Usually lasts' column. In children with middle-ear infection: if fluid is coming out of their ears or if they have deafness. Other
Sore throat	7 days		
Common cold	10 days		
Sinusitis	18 days		
Cough or bronchitis	21 days		
Other infection:	days		

Never share antibiotics and always return any unused antibiotics to a pharmacy for safe disposal















